Influence of Nurse Practitioner Practice Restrictions on Chronic Disease Disparities

Margo Brooks Carthon, PhD, RN, FAAN Tyson Family Term Chair in Gerontologic Research University of Pennsylvania School of Nursing Center for Health Outcomes & Policy Research





UNIVERSITY of PENNSYLVANIA SCHOOL of NURSING

Acknowledgements

- **Collaborators:** Heather Brom, Lusine Poghosyan, Jacqueline Nikpour, Jessie Chittams
- National Council of State Boards of Nursing Center for Regulatory Excellence (Brooks Carthon, Poghosyan, MPI)
 - Influence of Nurse Practitioner Practice Restrictions on Chronic Disease Health Disparities
- National Institutes of Nursing Research (R01NR014855; Aiken, PI)




Coronary artery disease (CAD) & disparities

Most prevalent heart disease in U.S.


Leading cause of death in the U.S.

Requires lifestyle modifications, routine cholesterol screening and interventions for dyslipidemia Age-adjusted death rates for heart disease, by race and Hispanic origin: 1999–2017



Source: Centers for Disease Control. (2019). *Health, United States Spotlight*. Racial and Ethnic Disparities in Heart Disease.

CAD disparities





PATIENT FACTORS

PROVIDER FACTORS

SETTINGS WITH FEWER RESOURCES

Majority of NPs deliver primary care

Nurse practitionerdelivered care

Numerous studies demonstrate that NPs deliver high quality care

Constraints to NP-delivered care

Restriction in scope of practice


Unsupportive practice environments


The NP Practice Environment

- Better practice environments

 - ↑ screening for cardiovascular disease







The NP Health Disparities Model (Poghosyan & Brooks Carthon, 2017)

Research question: Does the NP practice environment reduce racial disparities in LDL cholesterol screening between Black and White CAD patients?

Methods

Cross-sectional design

- RN4CAST-US NP survey (2015-2016)
- SK&A physician database (2016)
- Medicare claims (2016)


Setting & sample

- 456 primary care practices & 502 NPs across 4 states (CA, FL, NJ, PA)
- 111,911 CAD patients
 - 6% Black and 94% White

Data analysis


- Multivariable regression models
- Interaction term for race and practice environment

Variables, measures, and data sources




Results - patients

- 111,911 CAD patients
 6% Black and 94% White
- Black CAD patients were...
 - Younger (70 vs. 76 years, *p*<.001)
 - Female (58% vs. 47%, p<.001)
 - Enrolled in Medicaid (46% vs. 11%, p<.001)
- Average of 6 chronic conditions



Results - practices

- Practices where Black CAD patients received care...
 - Accepted Medicaid (73% vs. 59%, p<.001)
 - Smaller daily patient volume (166 vs. 193 patients/day, p<.001)
 - Less favorable practice environments (3.25 vs. 3.27, p<.001)




Results


Odds of LDL screening <u>19% lower</u> for Black vs. to White CAD patients


Significant interaction found between race and the practice environment

Probability of Lipid Testing for Black and White CAD Patients



Limitations





CROSS-SECTIONAL STUDY DESIGN

LDL SCREENING FROM BILLING DATA

Conclusions

The first study to examine the relationship between NP practice environments and health disparities.

Disparities in CAD management between Black and White patients may be influenced by where they receive care.

Investing in the NP practice environment, including supports for NP independent practice and strengthening NPs' relationships with physicians and administrators, may narrow these disparities.

Thank you!

Margo Brooks Carthon, PhD, RN, FAAN

jmbrooks@nursing.upenn.edu

